

The effect of supernova heating on cluster properties and constraints on galaxy formation models

K. K. S. Wu,^{1,3} A. C. Fabian¹ and P. E. J. Nulsen²

¹ *Institute of Astronomy, Madingley Road, Cambridge CB3 0HA*

² *Department of Physics, University of Wollongong, Wollongong NSW 2522, Australia*

³ *kwu@ast.cam.ac.uk*

ABSTRACT

Models of galaxy formation should be able to predict the properties of clusters of galaxies, in particular their gas fractions, metallicities, X-ray luminosity-temperature relation, temperature function and mass-deposition-rate function. Fitting these properties places important constraints on galaxy formation on all scales. By following gas processes in detail, our semi-analytic model (based on that of Nulsen & Fabian 1997) is the only such model able to predict all of the above cluster properties. We use realistic gas fractions and gas density profiles, and as required by observations we break the self-similarity of cluster structure by including supernova heating of intracluster gas, the amount of which is indicated by the observed metallicities. We also highlight the importance of the mass-deposition-rate function as an independent and very sensitive probe of cluster structure.

Key words: galaxies: clusters: general – galaxies: formation – galaxies: evolution – cooling flows – X-rays: galaxies

1 INTRODUCTION

Our understanding of galaxy formation through the hierarchical clustering of gas and dark matter (DM) has improved greatly over the past decade. The aim of models of galaxy formation is to account for the properties of all structure, from the first dwarf galaxy to the rich clusters of galaxies that are virialising now. The collapse of the dominant dark matter is relatively well understood, so that the major current problem is to model the complex gas processes that take place within collapsed and collapsing DM halos. Despite their success in accounting for some properties of galaxies, many semi-analytical models do not predict correctly the properties of groups and clusters of galaxies. In particular, these models should predict gas fractions, metallicities, the X-ray luminosity-temperature ($L_X - T$) relation, the temperature function and the mass-deposition-rate ($\dot{M}$) function of clusters.

One problem is that the gas fraction in clusters, which is observed to be in the range 10–20 per cent (White et al. 1993b; White & Fabian 1995; David, Jones & Forman 1995; White, Jones & Forman 1997; Evrard 1997), is underestimated by those models that have low baryon fractions (e.g., most of the values considered by Kauffmann, White & Guiderdoni (1993), Cole et al. (1994), Heyl et al. (1995), Kauffmann & Charlot (1998) and Baugh et al. (1998)).

In this paper we show that additional heat input to

the intracluster gas, which in our fiducial model we take to be due to supernovae, can consistently account for the major X-ray properties of groups and clusters of galaxies. This has a similar effect to ‘preheating’ as invoked by other authors (Kaiser 1991; Evrard & Henry 1991; Metzler & Evrard 1994; Navarro, Frenk & White 1995; Bower 1997; Jones et al. 1998). A well-known problem addressed by preheating is the slope of the $L_X - T$ relation. Models and simulations which do not involve preheating (and in general those which investigate galaxy properties do not) are generally ‘self-similar’ and give $L_X \propto T^2$ (Kaiser 1986), whereas clusters are observed to follow $L_X \propto T^3$ (David et al. 1993). Models which preheat the gas in clusters generally break the self-similarity. For example, a constant excess specific energy can have a large effect in the shallow potential wells of groups, while having little effect on the largest clusters. The excess energy raises the temperature as well as the potential energy of the gas by pushing it outwards, thereby flattening the gas density profile. This lowers L_X and steepens the relationship between L_X and T .

During galaxy formation gas is heated and possibly ejected from galaxies by supernovae. It retains most of this ‘excess energy’ as thermal and gravitational potential energy (and possibly kinetic energy) as it passes through the collapse hierarchy, eventually becoming part of the intergalactic medium in groups and clusters. Gas that is ejected from a galaxy is expected to recollapse with the next ‘major merger’

(which in our model is the doubling, at least, of the mass of the halo that the galaxy resides in). This is reasonable since ejected gas would encounter the inter-galactic medium which subsequently forms part of a larger halo. In our fiducial model the only source of heating and metal enrichment is Type II supernovae (SNe II) associated with star formation (Nulsen & Fabian 1995, 1997; hereafter NF95 and NF97). There is some evidence that most of the metals in intracluster gas came from SNe II rather than SNe Ia (Mushotzky et al. 1996; Nagataki & Sato 1998), but a further reason for our assumption is because while both types inject about the same amount of energy, SNe Ia inject about 10 times more iron than SNe II. Hence for the iron abundance measured in intracluster gas, the associated energy injection is maximised by using only SNe II. We assume an energy injection of 4×10^{50} erg per supernova (Spitzer 1978) which is not radiated locally. The mean mass of iron produced per supernova is $0.07 M_{\odot}$, which leads to 3.7 keV/particle excess specific energy in gas with solar iron abundance. By choosing the initial metallicity in groups when they are first formed to be a suitable value, the iron abundance of clusters in our model is $Z \approx 0.3Z_{\odot}$ which corresponds to about 1.1 keV/particle of excess energy.

The semi-analytical model we use is based on that described in NF97, with an improved cosmological model and a more accurate form for dark matter halos. The model pays particular attention to gas processes, explicitly including the effects of cooling flows in its treatment of hot gas. When hot gas has had sufficient time to radiate its thermal energy it is assumed to preferentially form low mass stars, effectively baryonic dark matter (BDM), as observed in cooling flows in clusters. We assume that dark matter halos follow the NFW profile (Navarro, Frenk & White 1997; hereafter NFW97). Gas in the resulting potential wells is assumed to be isothermal and in hydrostatic equilibrium. The model clusters turn out to have gas density profiles closely resembling those in observed clusters.

In this paper we use an open cosmology with $\Omega_0 = 0.3$, $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and a cosmological constant of zero. Density fluctuations are obtained from a CDM model with primordial spectral index of $n = 1$, normalised to make $\sigma_8 = 0.7$. This normalisation, which is slightly lower than some estimates based on cluster numbers, is discussed in section 3. We note that our results would be largely unchanged in a flat cosmology with the same Ω_0 , since the mass function of clusters today would be the same. Likewise, our conclusions do not depend on when the excess energies were injected or the exact history of the clusters and their progenitors. We do however assume that the supernovae injected their energy into virialised gas, as opposed to the uncollapsed IGM.

2 DESCRIPTION OF THE MODEL

The two main changes from the model described in NF97 are the use of an open cosmology and the improvement of the DM and gas profiles. The mass of a halo is defined as that inside the virial radius, r_{200} , and the mean density inside r_{200} is required to be 200 times the background density of an $\Omega = 1$ universe of the same age. (This is marginally different from NFW97, but follows the spherical collapse model strictly.) We use the NFW profile (NFW97) which takes the form

$$\rho(r) = \frac{\rho_s}{x(1+x)^2}, \quad (1)$$

where $x = r/r_s$, and the characteristic density and scale, ρ_s and r_s , are parameters. For a given mass virialising at some time in a given cosmology (which yields r_{200}), $\rho(r)$ has only one degree of freedom, which can be fixed by applying the algorithm given in the Appendix of NFW97 to obtain the concentration, $c = r_{200}/r_s$. ρ_s is then fixed since we know the mean density inside r_{200} . The algorithm used was shown by NFW97 to provide reasonable fits of c to simulation results, where small halos were found to be more ‘concentrated’ than large ones. In our cosmology the value of c typically ranges from 4.5–6.5 for clusters.

We consider the above profile to describe the total density of DM and gas, thus the gravitational potential $\phi(r)$ is calculated from it, assuming that the halo is truncated at r_{200} :

$$\phi(r) = \alpha \left(-\frac{\ln(1+x)}{x} + \frac{1}{1+c} \right), \quad (2)$$

where $\alpha = 4\pi G \rho_s r_s^2$. Since the gas fraction is small, the DM also follows the NFW profile to a good approximation.

In our model the gas is isothermal and in hydrostatic equilibrium, giving the gas density profile

$$\rho_g(r) \propto \exp \left(-\frac{\mu m_H}{kT} \phi(r) \right), \quad (3)$$

where μm_H is the mean mass per particle in the gas, T is its temperature and k is Boltzmann’s constant. Inserting the expression for the potential gives

$$\rho_g(r) = A(1+x)^{\eta/x} \quad (4)$$

where A is a constant and $\eta = \mu m_H \alpha / (kT)$. The above profile, which we shall refer to as the NFW-gas profile, was shown by Makino et al. (1997) to closely approximate the conventional β -model (with $\beta = \eta/15$). It tends to Ae^{η} in the centre and to A for large x . Thus A is, in practice, very small and the plateau at large x is not noticeable. The parameter α has the same units as velocity dispersion squared, thus it is apparent that η is closely related to the widely-used parameter β , the ratio of the DM kinetic energy to the thermal energy of the gas when both are isothermal.

The NFW-gas profile has been fitted to the surface brightness profiles of observed clusters and it models most clusters very well (Ettori & Fabian 1998). Values of η in the largest clusters which have cooling flows (suggesting they are in a relaxed state, in contrast to non-cooling flows; Buote & Tsai 1996) average about 10.5, with a standard deviation of about 1.0 (Ettori, private communication). Direct comparison shows that the gas profile for $\eta = 10.5$ closely follows the NFW profile from $x = 0.4$ to 3. The profiles differ inside $x = 0.4$ due to the flat core of the NFW-gas profile, which implies that the gas fraction increases as a function of radius, as observed in real clusters (e.g. White, Jones & Forman 1997). This last effect becomes more pronounced and spreads to larger radii for smaller values of η , the major cause of which will be heating by supernovae.

Since η gives both the gas temperature and the slope of the gas density profile, it is the structural parameter through which supernova heating exerts its influence on cluster properties. The procedure for setting η is as follows. We postulate that the total specific energy of the gas (thermal plus gravi-

tational) is proportional to the specific gravitational energy of the DM, and that any excess specific energy from supernova heating is added to the above gas energy, as its name implies. i.e.,

$$\frac{3kT}{2\mu m_H} + \frac{\int \rho_g \phi dV}{M_{\text{gas}}} = K \frac{\int \frac{1}{2} \rho_{\text{DM}} \phi dV}{M_{\text{DM}}} + E_{\text{excess}}, \quad (5)$$

where ρ_{DM} is the DM density, which follows the NFW profile, M_{gas} and M_{DM} are the total gas and DM masses, and E_{excess} is the excess specific energy. The volumes of integration are spheres of radius r_{200} . Given the constant K , it is easy to show that η is only a function of c and E_{excess} . The constant of proportionality K is calibrated so that η is close to the observed value of 10.5 for the largest clusters, when $E_{\text{excess}} = 0$. We use the value $K = 1.2$. (Note that η varies by around 0.5 depending on collapse redshift, through its dependence on c .) The reason for matching to the largest clusters is that if supernova heating does occur, we expect it to have least effect on the largest clusters. As c varies little over cluster masses, this method ensures that clusters are self-similar when there is no excess energy (i.e., c and η are roughly constant). The trends for c and η are to increase slightly with decreasing mass, over all masses in our model (see below).

The mass deposition rates of clusters are estimated by finding the radius r_{cool} at which the cooling time is equal to the time Δt to the next major merger (see below) or the present, and dividing the gas mass enclosed by Δt to give $\dot{M}$. For clusters with cooling flows, i.e., those with significant $\dot{M}$, the hot gas which cools does not form stars but BDM.

The Cole & Kaiser (1988) block model is used to simulate the merging history of dark matter halos. The smallest masses in the hierarchy are $1.5 \times 10^{10} M_{\odot}$ and there are 20 levels of hierarchy so that the total mass in one collapse tree is $7.9 \times 10^{15} M_{\odot}$. When a ‘block’ in the hierarchy virialises, it merges into one halo both older halos and previously uncollapsed material. The new average metallicity and excess energy of the gas can therefore be calculated. Since blocks double in mass up the hierarchy, each collapse of a block roughly corresponds to a major merger. As already mentioned, we set the metallicities in groups to be high without considering the implications of this for star formation. Halos with masses in the range $15\text{--}120 \times 10^{12} M_{\odot}$ are given initial iron abundances of $Z = 0.5Z_{\odot}$ (This implies that the gas in some of the smaller halos is unbound.) As a result the average metallicities (for a given mass) range from $0.3Z_{\odot}$ for clusters of mass $250 \times 10^{12} M_{\odot}$ to $0.25Z_{\odot}$ for the largest clusters. The range of scatter in metallicities is $\approx 0.1Z_{\odot}$. These are in good agreement with the recent *ASCA* measurements by Fukazawa et al. (1998) of iron abundances in 40 nearby clusters and the measurements of Allen & Fabian (1998).

The gas fraction of the clusters in our model lie within one per cent of 14 per cent.

3 RESULTS AND DISCUSSION

To illustrate the effect of heating from supernovae, we display in Fig. 1 the L_X - T distributions before and after the inclusion of supernova heating. We superimpose on the distributions the power law fit obtained by David et al. (1993), for bolometric luminosities. The excess energies bring the




Figure 1. Contour plots of the luminosity-temperature distribution, without heating by supernovae (dotted lines) and with supernova heating (solid lines). The contours are spaced at equal logarithmic intervals. The long straight line is the best fit (for bolometric luminosities) taken from David et al. (1993). The distribution obtained without heating is roughly parallel to the short line segment which follows $L_X \propto T^2$. In this case the smallest clusters are overluminous by an order of magnitude. The excess energies bring the smaller clusters into agreement with data.

smaller clusters into agreement with the observed correlation, by increasing their temperature, flattening their density profiles and thereby decreasing their luminosities. For clusters of mass $2.5 \times 10^{14} M_{\odot}$, η decreases to around 9. The spread in the L_X - T distributions are significant and arise naturally in our simulation from the different formation histories of the clusters.

For comparison with Fig. 1, if we halve the excess energies injected, the average luminosity of a 2×10^7 K cluster rises to $5 \times 10^{43} \text{ erg s}^{-1}$. Doubling the excess energies causes most of the clusters with $T < 3 \times 10^7$ K to be unbound, which clearly disagrees with data, and roughly halves the luminosity of 3×10^7 K clusters. In future work we will investigate whether restricting the heating to the core of clusters can decrease the excess energies required (preliminary results suggest that this could reduce the total energy requirements by about a third).

In Figs. 2 and 3 we show the X-ray luminosity function (XLF) and temperature function respectively. The curves plotted for comparison are the best fit Schechter function to the *ROSAT* Brightest Cluster Sample (BCS) bolometric luminosity function, obtained by Ebeling et al. (1997), and the best power law fit to the temperature function obtained by Edge et al. (1990). The errors in the BCS XLF are very small, while the temperature function is currently uncertain by factors of around 2 or 3. A problem which follows from an incorrect slope of the L_X - T relation is that if the temperature function fits the data reasonably well, then the XLF is too steep compared to data (e.g. Kitayama and Suto 1996). Correcting the L_X - T relation thus improves the slope of the XLF significantly, though it is still slightly steeper than observed.

Estimates of σ_8 from cluster abundances in general use a theoretically derived mass function and assume some mass-




Figure 2. The X-ray luminosity function (XLF) from the model with supernova heating. The curve is the best fit Schechter function to the *ROSAT* Brightest Cluster Sample (BCS) bolometric luminosity function (Ebeling et al. 1997).




Figure 3. The temperature function from the model with supernova heating. The power law fit to the temperature function obtained by Edge et al. (1990) is plotted.

temperature ($M - T$) relation in order to compare with an observed temperature function. The calibration of the $M - T$ relation is thus crucial to the results, especially as the temperature function has a very steep slope. We find that with zero excess energy our model clusters obey the simple estimate (appropriate for isothermal r^{-2} gas and DM density profiles) of

$$\frac{GM_{200}}{2r_{200}} = \frac{kT}{\mu m_H} \quad (6)$$

to within a few per cent, where M_{200} is the total mass within r_{200} . Supernova heating raises the temperature slightly for small clusters. For a cluster of mass $4.9 \times 10^{14} M_\odot$ collapsing at $z = 0$ the temperature increases from 3.5×10^7 K to 3.8×10^7 K in the model described, close to 10 per cent higher. The resulting temperatures in our model are a necessary




Figure 4. The smoother curve is the $\dot{M}$ cumulative function from our model with supernova heating, and is compared to the function taken from Peres et al. (1997) but with the age of clusters changed to 6 Gyr, as explained in the text. We have also excluded clusters with $T < 2 \times 10^7$ K; when all clusters are allowed to contribute, the function increases below $50 M_\odot \text{ yr}^{-1}$, rising to $6 \times 10^{-5} \text{ Mpc}^{-3}$ at $10 M_\odot \text{ yr}^{-1}$, but such clusters would not be included in the observed sample.

consequence of fitting the $L_X - T$ relation. In a recent paper, Eke et al. (1998) used a more elaborate form of Equation 6 and included scatter in the $M - T$ relation, which brought down the predicted σ_8 . With our CDM shape parameter of $\Gamma = \Omega_0 h \exp(-\Omega_b(1 + 1/\Omega_0)) = 0.12$ their Fig. 7 predicts a value of $\sigma_8 \approx 0.75$. Since preheating was not included, our value of $\sigma_8 = 0.7$ appears consistent with their results.

Turning to the mass deposition rates, we display the $\dot{M}$ cumulative space density in Fig. 4. We have excluded clusters with temperatures less than 2×10^7 K since such objects would be too faint in the 2–10 keV band to be included in the sample used by the two papers below. Our result agrees very well with the $\dot{M}$ cumulative function obtained by Edge, Stewart & Fabian (1992) and lies slightly lower than that shown in Peres et al. (1997). The systematic uncertainties in measurements of $\dot{M}$ in clusters are at present around a factor of 2. In particular the age of a cluster since its last major merger is unknown and in determining r_{cool} the age of the universe is generally used. Lowering this age in the analysis of Peres et al. (1997) from 13 Gyr to 6 Gyr, which is the typical age of a 5 keV cluster in our model, lowers r_{cool} and more than halves the values of $\dot{M}$ in several clusters (Peres, private communication). In Fig. 4 we show the cumulative function from Peres et al. (1997) recalculated with this change in the age of clusters. There is very good agreement for all but small values of $\dot{M}$, which is not a serious concern since the clusters and their measurements are both sensitive to small perturbations in this range.

The mass deposition rate of a cluster is a highly sensitive probe of the gas density inside a radius of ~ 0.1 Mpc (the typical r_{cool} observed in cooling flows). On the other hand, the luminosity in all but the most massive cooling flows comes from inside a radius $\sim r_s$ which is larger than the cooling flow region. Therefore L_X and $\dot{M}$ are independent probes of the gas density profile at different radii, and due to their sensitivity they are efficient at eliminating mod-

els. For example, while we can adjust the overall density inside $\sim r_s$ by changing the temperature, the ratio of the densities inside the two radii described above have to be correct in order to match both the $\dot{M}$ function and the XLF. We found that models which used gas density profiles of the form $r^{-2\beta}$ (see NF97) overpredicted the $\dot{M}$ function by an order of magnitude or more. The agreement with the available data suggests that the NFW-gas profile is much closer to the right shape.

As a further example of the ability of $\dot{M}$ to discriminate between models, without supernova heating we find an average value of $\dot{M} \approx 100 M_\odot \text{yr}^{-1}$ for cooling flow clusters with $L_X = 6 \times 10^{43} \text{erg s}^{-1}$ which is too high when compared with Fig. 2 of Peres et al. (1997). However, heating lowers the value to $\dot{M} \approx 30 M_\odot \text{yr}^{-1}$, which then agrees with their data. On the other hand, large clusters with $L_X = 10^{45} \text{erg s}^{-1}$ have the correct $\dot{M}$ with and without heating. Using a hydrodynamic and N-body simulation, Katz & White (1993) modelled a Virgo-like cluster with cooling of gas included, and obtained an excessively high value of $\dot{M} \sim 400 M_\odot \text{yr}^{-1}$. Our results suggest that non-gravitational heating could simultaneously help solve such a problem. The above results argue for flatter gas density profiles in small clusters, independently of the $L_X - T$ relation. We consider it very promising that by correctly fitting the $L_X - T$ relation, the predictions pertaining to $\dot{M}$ naturally agree with the data.

4 CONCLUSIONS

We have presented a galaxy formation model in which the X-ray properties of clusters agree well with what is observed. The model is based on that described by Nulsen & Fabian (1997), and uses observed gas fractions and metallicities. Realistic gas density profiles were used, which sit in potentials given by the NFW profile. In particular, we obtained good fits to the $L_X - T$ relation, the X-ray luminosity function, the temperature function and the $\dot{M}$ function, which we find to be a very sensitive diagnostic of cluster structure in the core.

To resolve a number of problems, including getting the correct slope for the $L_X - T$ relation and avoiding excessive mass deposition rates for low luminosity clusters, self-similarity of clusters has to be broken. In our model this is achieved by flattening the gas density profiles of the smaller clusters, which occurs naturally when we include heating of intracluster gas due to the retention of energy from supernovae in all previous collapse stages. We find that the energy required agrees well with the supernova energy released in producing the observed metallicities if most of this energy is retained in the intracluster gas. The result is that the above problems are simultaneously resolved once supernova heating is included.

We have highlighted the importance of the mass deposition rate and the $\dot{M}$ function as diagnostics of cluster structure. The value of $\dot{M}$ probes the gas density profile at smaller radii than L_X and hence is able to give us independent information. Due to the sensitivity of quantities such as L_X and $\dot{M}$ to the structure and density of the hot gas, and the steep slopes of the three distribution functions we consider, detailed gas processes can be as important as cosmological considerations in determining what we observe.

ACKNOWLEDGEMENTS

We thank Stefano Ettori for the results of surface brightness profile fits, Clovis Peres for the new calculation of the $\dot{M}$ function, and Vince Eke, Stefano, Clovis and Martin Rees for very helpful discussions. KKS is grateful to the Croucher Foundation for financial support. ACF thanks the Royal Society for support. PEJN gratefully acknowledges the hospitality of the Institute of Astronomy during part of this work.

REFERENCES

- Allen S. W., Fabian A. C., 1998, MNRAS, 297, L63
- Baugh C. M., Cole S., Frenk C. S., Lacey C. G., 1998, ApJ, 498, 504
- Bower R. G., 1997, MNRAS, 288, 355
- Buote D. A., Tsai J. C., 1996, ApJ, 458, 27
- Cole S., Aragón-Salamanca A., Frenk C. S., Navarro J. F., Zepf S. E., 1994, MNRAS, 271, 781
- Cole S., Kaiser N., 1988, MNRAS, 233, 637
- David L. P., Jones C., Forman W., 1995, ApJ, 445, 578
- David L. P., Slyz A., Jones C., Forman W., Vrtilek S. D., 1993, ApJ, 412, 479
- Ebeling H., Edge A. C., Fabian A. C., Allen S. W., Crawford C. S., 1997, ApJ, 479, L101
- Edge A. C., Stewart G. C., Fabian A. C., 1992, MNRAS, 258, 177
- Edge A. C., Stewart G. C., Fabian A. C., Arnaud K. A., 1990, MNRAS, 245, 559
- Eke V. R., Cole S., Frenk C. S., Henry J. P., 1998, MNRAS, submitted (astro-ph/9802350)
- Ettori S., Fabian A. C., 1998, MNRAS, submitted
- Evrard A. E., 1997, MNRAS, 292, 289
- Evrard A. E., Henry J. P., 1991, ApJ, 383, 95
- Fukazawa Y., Makishima K., Tamura T., Ezawa H., Xu H., Ikebe Y., Kikuchi K., Ohashi T., 1998, PASJ, 50, 187
- Heyl J. S., Cole S., Frenk C. S., Navarro J. F., 1995, MNRAS, 274, 755
- Jones L. R., Scharf C., Ebeling H., Perlman E., Wegner G., Malkan M., Horner D., 1998, ApJ, 495, 100
- Kaiser N., 1986, MNRAS, 222, 323
- Kaiser N., 1991, ApJ, 383, 104
- Katz N., White S. D. M., 1993, ApJ, 412, 455
- Kauffmann G., Charlot S., 1998, MNRAS, 294, 705
- Kauffmann G., White S. D. M., Guiderdoni B., 1993, MNRAS, 264, 201
- Kitayama T., Suto Y., 1996, ApJ, 469, 480
- Makino N., Sasaki S., Suto Y., 1997, ApJ, 497, 555
- Metzler C. A., Evrard A. E., 1994, ApJ, 437, 564
- Mushotzky R., Loewenstein M., Arnaud K. A., Tamura T., Fukazawa Y., Matsushita K., Kikuchi K., Hatsukade I., 1996, ApJ, 466, 686
- Nagataki S., Sato K., 1998, ApJ, in press (astro-ph/9807022)
- Navarro J. F., Frenk C. S., White S. D. M., 1995, MNRAS, 275, 720
- Navarro J. F., Frenk C. S., White S. D. M., 1997, ApJ, 490, 493 (NFW97)
- Nulsen P. E. J., Fabian A. C., 1995, MNRAS, 277, 561 (NF95)
- Nulsen P. E. J., Fabian A. C., 1997, MNRAS, 291, 425 (NF97)
- Peres C. B., Fabian A. C., Edge A. C., Allen S. W., Johnstone R. M., White D. A., 1998, MNRAS, in press (astro-ph/9805122)
- Spitzer L., 1978, Physical Processes in the Interstellar Medium. Wiley-Interscience, New York
- White D. A., Fabian A. C., 1995, MNRAS, 273, 72
- White D. A., Jones C., Forman W., 1997, MNRAS, 292, 419
- White S. D. M., Navarro J. F., Evrard A. E., Frenk C. S., 1993, Nature, 366, 429